arXiv:1508.00195v1 [math.NT] 2 Aug 2015

Dimension groups and linear diophantine inequalities

Abstract Solvability of systems of linear diophantine inequalities is related to dimension group properties, yielding necessary and sufficient conditions. Specifically, if H is a subgroup of Euclidean space, for every h in H and positive integer m, for every $\epsilon > 0$, we can solve for x in H the inequalities $h - mx > -\epsilon$ coordinatewise, iff for F defined as the smallest face of the standard simplex (in the dual of the Euclidean space) that contains the positive linear functionals killing H, for all f in F, the subgroup of the reals, $f(H)$, is either zero or dense.

David Handelman & Damien Roy[1]

Introduction

This paper answers some questions in [BeH] concerning refinable measures and more generally unperforation of quotients by (some) subgroups, and at the same time, yields results about classes of diophantine inequalities. Here is a motivating example for the latter.

EXAMPLE 1 Let A be a real square matrix of size $n-2$. Define the $(n-1)\times n$ real matrix,

$$B = \begin{pmatrix} \alpha_1 & -\alpha_1 & & & & & & \\ \alpha_2 & -\alpha_2 & & & & & & \\ \vdots & \vdots & & & A & & & \\ \alpha_{n-2} & -\alpha_{n-2} & & & & & & \\ \alpha_{n-1} & -\alpha_{n-1} & \eta_1 & \eta_2 & \eta_3 & \dots & \eta_{n-3} & \eta_{n-2} \end{pmatrix}.$$

Denote the rows w_i. Suppose that $\sum \alpha_i \mathbf{Z}$ has rank at least two (that is, at least one of the ratios α_i/α_j is irrational). Let H be the abelian group generated by $\{w_i\}$, that is, the row space of B. We ask whether H satisfies the following property (whose origin will be explained later).

(*) For all $h \in H$ and $\epsilon > 0$, there exists $h' \in H$ (depending on ϵ and h) such that all coordinates of $h - 2h'$ exceed $-\epsilon$.

The 2 that appears in (*) can be replaced by any positive integer $m > 1$ (these are all equivalent). Applying this in the case that $A = \mathrm{I}$ and with $h = v_1$, the property asks whether given $\epsilon > 0$ and $m > 1$, there exist integers $a(i)$ such that the inequalities

$$\left|\frac{\alpha_1}{m} - \sum a(i)\alpha_i\right| < \epsilon$$
$$\frac{1}{m} - a(1) - a(n-1)\eta_1 \geq -\epsilon$$
$$-a(j) - a(n-1)\eta_j \geq -\epsilon \quad \text{for all } 1 < j < n-1$$

hold. (The inequalities derive from considering $h/m - h'$ and replacing ϵ by $m\epsilon$.) It is possible to prove by rather indirect methods that (*) holds when $\{w_i\}$ is real linearly independent, while if $\{w_i\}$ is not real linearly independent, then (*) holds if $\{1, \eta_1, \dots, \eta_{n-2}\}$ is rationally linearly independent. Our principal result will give this quickly, and much more generally (for example, if A is *any* nonsingular matrix and the rows of B form a linearly independent set, then (*) still holds).

With $A = \mathrm{I}$ but deleting the last row, (*) fails, meaning that for at least one choice of $h = w_i$, there exists $\delta > 0$ such that $h - 2h'$ has a negative entry of magnitude at least δ for all $h' \in H$.

Another simple class of examples, depending on r, are of the form,

$$\begin{pmatrix} 1 & -1 & 1 & 0 \\ \alpha & -\alpha & r & 1 \end{pmatrix},$$

---

[1] Both authors supported in part by NSERC Discovery grants.

Keywords and phrases: diophantine inequalities, dimension group, trace, Gordan's Theorem, unperforation, refinable measure, simplex

AMS (MOS) 2010 classification: 19K14 (primary); 11J25, 15A39, 06F20, 37A55

where α is irrational. These are easy to analyze, but display some interesting dependence on r.

The main result with respect to diophantine inequalities, described in the abstract, reduces the existence of solutions to determining the behaviour of *positive* linear functionals on H.

Dimension group preliminaries

Let G be a partially ordered abelian group with positive cone denoted G^+; we include *directedness* in the definition, that is, $G = G^+ - G^+$. An *order unit* (or *strong unit*) of G is an element, u, of G such that for all $g \in G$, there exists a positive integer N with $-Nu \leq g \leq Nu$. For example, there two obvious partial ordering on $\mathbf{R}^n$, the usual coordinatewise one (with positive cone $(\mathbf{R}^n)^+$ consisting of vectors all of whose entries are nonnegative), and the *strict ordering,* whose positive cone consists of zero, together with the strictly positive vectors (all entries strictly greater than zero). The order units of $\mathbf{R}^n$ with respect to either ordering are the strictly positive vectors. The set of order units of a partially ordered group G will be denoted G^{++}.

We adopt the convention that if (G, u) is a partially ordered abelian group with order unit, then $u \neq 0$ (so that G is nontrivial).

A partially ordered group G is *unperforated* if whenever n is a positive integer and $g \in G$, $ng \in G^+$ implies $g \in G^+$. The orderings on $\mathbf{R}^n$ described in the previous paragraph are both unperforated.

A *trace* (or *state*) on G is a nonzero group homomorphism $\tau : G \to \mathbf{R}$ that is positive (meaning $\tau(G^+) \subset \mathbf{R}^+$). If u is an order unit for G, the trace τ is *normalized at* u if $\tau(u) = 1$. On $\mathbf{R}^n$ with either the usual or the strict ordering, the traces are given by nonnegative linear combinations of the coordinate functions, denoted τ_i, that pick out the ith coordinate. If we set $u = (1, 1, \ldots, 1)$, so it is an order unit in either ordering, the normalized traces constitute the standard simplex (in the dual space),

$$\left\{ \sum \lambda_i \tau_i \,\middle|\, \lambda_i \geq 0,\ \sum \lambda_i = 1 \right\}.$$

A *dimension group* is an unperforated partially ordered abelian group that satisfies the *Riesz decomposition property,* that is, whenever $a \leq b + c$ with $a, b, c \in G^+$, then there exist $0 \leq b_1 \leq b$ and $0 \leq c_1 \leq c$ such that $a = b_1 + c_1$. Both orderings on $\mathbf{R}^n$ satisfy this. Dimension groups are precisely the direct limits of *simplicial groups* ($\mathbf{Z}^n$ with the usual ordering). A partially ordered abelian group is *simple* if every nonzero element of the positive cone is an order unit. Thus $\mathbf{R}^n$ in the strict ordering is simple, but if $n > 1$, $\mathbf{R}^n$ is not simple in the usual ordering. In particular, all simple partially ordered abelian groups admit order units, and in that case, we have $G^{++} = G^+ \setminus \{0\}$. We will mostly be concerned with simple dimension groups.

With (G, u) a partially ordered group with order unit, we can form $S(G, u)$, the set of normalized traces on G; with respect to the weak topology, this is a compact convex set. When G is a dimension group, $S(G, u)$ is a Choquet simplex. The affine representation of (G, u) is essentially the second dual map, $\widehat{\ } : G \to \operatorname{Aff} S(G, u)$ given by $g \mapsto \widehat{g}$, where $\widehat{g}(\tau) = \tau(g)$. This is an order-preserving group homomorphism (where $\operatorname{Aff} S(G, u)$ is equipped with the usual, coordinatewise ordering).

For more on dimension groups and their affine representations, see [G].

Connections between minimal $\mathbf{Z}$-actions on Cantor sets and simple dimension groups led [BeH] to consider quotients of the following sort.

Let G be a partially ordered abelian group, and H a subgroup of G such that $H \cap G^+ = \{0\}$. On G/H, we can define a pre-ordering, $(G/H)^+ = \{g + H \mid \exists h \in H,\ g + h \in G^+\}$. This is a partial ordering, and the quotient group will be simple, if G is. Now assume G admits an order unit, u; then $u + H$ is an order unit for G/H (since $H \cap G^+ = \{0\}$ and G is directed, $H \neq G$). This is a special case of a more general result about convex subgroups, which we will discuss later.

The traces on G/H are precisely those of G that kill H. In general of course, G/H need not be even torsion-free, let alone unperforated. Suppose $U \subset S(G, u)$ (or U could be just a set of traces, not necessarily normalized), and define $\ker U = \ker{}_G U = \{g \in G \mid \tau(g) = 0\, \forall\, \tau \in U\}$. Where the choice of G is unambiguous, we may drop the subscript G. Then $G/\ker{}_G U$ is obviously torsion-free, but need not be unperforated. There is a simple criterion for G/H to be unperforated when G is a simple dimension group. The following includes the converse to [BeH, B1] (which was left open there).

PROPOSITION 1.1 Let (G, u) be a simple dimension group, and let H be a subgroup such that $G^+ \cap H = \{0\}$ and G/H is torsion-free. Then the partially ordered group G/H is unperforated iff

(*) for all $h \in H$, for all $\epsilon > 0$, there exists $h' \in H$ such that $\widehat{h} - 2\widehat{h'} \geq -\epsilon$.

Proof. Sufficiency is proved in [BeH, B1]. For necessity, we may assume that G is not cyclic, so by [GH2, 4.9], the image of G under the affine representation map $\widehat{\ }$, is norm dense in $\operatorname{Aff} S(G,u)$. Given $h \in H$, there thus exists $g' \in G$ such that $\|\widehat{g'} - \widehat{h}/2\| < \epsilon/6$. Again by density, there exists $g'' \in G$ such that $\epsilon/6 < \widehat{g''} < 2\epsilon/5$. Thus with $g = g' + g''$, we have $3\epsilon/6 \gg \widehat{g} - \widehat{h}/2 \gg 0$. The right inequality means that $\widehat{2g-h} \gg 0$; by unperforation of G [GH1], this implies $2g - h \in G^+$. In particular, $2(g+H) \in (G/H)^+$. Unperforation of G/H yields $g + H \in (G/H)^+$.

Thus there exists h'' in H such that $g + h'' \in G^+$. From $2\widehat{g} + \widehat{h''} + (6\epsilon/6) - 2\widehat{g} - \widehat{h} \gg 0$, we deduce $2\widehat{h''} + \widehat{h} \gg -\epsilon$. Set $h' = -h''$. •

(The proof of the converse only requires density of the image of G in its affine representation together with unperforation of G, not the stronger requirement that it be a simple dimension group.)

The kernel of the map $G \to \operatorname{Aff} S(G,u)$ is known as the *infinitesimal subgroup* of G, denoted $\operatorname{Inf} G$. When G is a simple dimension group (other than $\mathbf{Z}$), the image of G is dense in $\operatorname{Aff} S(G,u)$, and moreover, G^+ consists of the pre-image of the strictly positive elements (the order units) of $\operatorname{Aff} S(G,u)$, together with zero.

Conversely, if G is a torsion-free abelian group and there is a group homomorphism $\phi : G \to \operatorname{Aff} K$ such that K is a Choquet simplex and the range of ϕ is norm dense, then the positive cone defined by $\phi^{-1}\left(\operatorname{Aff} K^{++}\right) \cup \{0\}$ will make G into a simple dimension group. We will be restricting ourselves to the situation in which K is finite-dimensional; equivalently, $\phi(G)$ is a dense subgroup of $\mathbf{R}^n$, and the ordering is the relative ordering from $\mathbf{R}^n$ equipped with the strict ordering.

Now suppose that (G,u) is a simple dimension group with n extreme (pure) traces; then $S(G,u)$ is just an $(n-1)$-simplex (a cross-section of the standard positive cone in $\mathbf{R}^n$), and $\operatorname{Aff} S(G,u) \cong \mathbf{R}^n$ as partially ordered vector spaces (the latter with the coordinatewise ordering).

Let H be a subgroup of G, fix an order unit u of the latter, and define

$$Z(H) = \{\tau \in S(G,u) \mid \tau(h) = 0 \text{ for all } h \in H\}.$$

This is the intersection of $S(G,u)$ with a closed flat, and is a compact convex subset of $S(G,u)$. For a set U of traces (or equivalently, a subset of $S(G,u)$), recall the definition, $\ker U = \{g \in G \mid \tau(g) = 0 \text{ for all } \tau \in U\}$ (this obviously depends on the choice of G; where there may be ambiguity, we write $\ker_G U$). It is routine to check that if $H = \ker_G U$ for some U, then $H = \ker_G Z(H)$. (Notice that $Z(H)$ only depends on the (the image of) H in $\mathbf{R}^n = \operatorname{Aff} S(G,u)$, while $\ker_G$ obviously depends on G.)

Recall from [BeH], that if $U \subset S(G,u)$, then the affine representation $G \to \operatorname{Aff} S(G,u)$ ($g \mapsto \widehat{g}$ where $\widehat{g}(\phi) = \phi(g)$) maps $\ker U$ into $U^{\vdash} := \{h \in \operatorname{Aff} S(G,u) \mid h|U \equiv 0\}$. Set J to be the closure the image of $\ker U$ (in $\operatorname{Aff} S(G,u)$), and define $P(J)$ as the subgroup of J generated by $\{h \in J \mid h \geq 0\}$. When $P(J)$ is dense in J, then [BeH, B2] applies. Our first automatic unperforation result is Theorem 1.2.

For a face F of a Choquet simplex, there exists a unique complementary face F' ($F \cap F' = \emptyset$, but $F \vee F' = S(G,u)$). If the Choquet simplex is finite-dimensional, then F' is closed and its extreme points are the extreme points of $S(G,u)$ not in F.

THEOREM 1.2 Suppose $Z(U)$ is a face whose complementary face is closed. Then $P(J) = J$, where J is the closure of the image of $\ker U$ in $\operatorname{Aff} S(G,u)$. In particular, $G/\ker U$ is unperforated; if G is additionally simple, then $G/\ker U$ is a dimension group.

Remark. Obviously $Z(U) := F$ is closed, and as $S(G,u)$ is a Choquet simplex, the closed face F has a (necessarily unique) complementary face, F'. The condition that F' be closed is unnecessary when $S(G,u)$ is finite-dimensional (that is, when it is an ordinary simplex in Euclidean space), but very restrictive when $S(G,u)$ is infinite-dimensional. For example, if X is a compact connected Hausdorff space, then $C(X,\mathbf{R}) = \operatorname{Aff} S(\mathcal{M}_1)$ (where $\mathcal{M}_1$ is the Bauer simplex consisting of Borel probability measures on X). The complementary face of any proper closed face is not closed (closed faces are in bijection with closed subsets of X).

Remark. Care should be taken with the hypotheses: they require that $Z(\ker_G U)$ be a face. This is a fairly strong condition; for example, it is easy to construct a pure trace τ on a simple dimension group such that $Z(\ker \tau)$ is not a face (while $\{\tau\}$ is a face), for which the conclusion fails.

We require a preparatory lemma. Lemma 1.3 is a generalization of Gordan's lemma, a version of Farkas' lemma concerning nonnegative solutions of linear systems of equations. This will be discussed in slightly more detail in the Appendix.

A subgroup H of a partially ordered abelian group G is *convex* if $h_1 \leq g \leq h_2$ with $h_i \in H$ and $g \in G$ entails that $g \in H$. If H is a convex subgroup unequal to G, then it is a routine to verify that G/H, equipped with the pre-ordering given by $(G/H)^+ = \{g + H \mid (g+H) \cap G^+ \neq \emptyset\}$, is partially ordered (that is, $(G/H)^+ \cap -(G/H)^+ = \{0 + H\}$, and $(G/H)^+$ generates G/H as an abelian group [G; Lemma 1.11]); moreover, if G admits an order unit, then so does G/H.

LEMMA 1.3 Let (G,u) be a partially ordered abelian group with order unit, and let L be a subgroup thereof. The following are equivalent.

(i) There exists a trace t of G such that $t(L) = \{0\}$;

(ii) $L \cap G^{++} = \{0\}$.

Proof. (i) implies (ii) is trivial, since order units are strictly positive at every trace.

(ii) implies (i). Set $c(L) = \{g \in G \mid \exists j_1, j_2 \in L \text{ such that } j_1 \leq g \leq j_2\}$. Then $c(L)$ is a convex subgroup of G; this follows immediately from the definitions, as does $L \subset c(L)$, and that $c(L)$ contains no order units of G. In particular, $c(L)$ is proper and convex. By the comments preceding this, $(G/c(L), u + c(L))$ is a partially ordered abelian group with order unit. By [GH, 3.3] or [G, Corollary 4.4], there exists a trace on $(G/c(L), u + c(L))$. Therefore, the composed map $G \to G/c(L) \to \mathbf{R}$ is a trace on G killing $c(L)$ and thus killing L. •

Remark. The group $c(L)$ constructed in the proof is the smallest convex subgroup of G containing L (c stands for *convex*), and the point underlying the argument is that $L \cap G^{++} = \emptyset$ implies that $c(L) \cap G^{++} = \emptyset$.

Proof. (of theorem 1.2) We have $Z(\ker U) = F$ is a closed face; let F' denote its complementary face, which we have assumed is closed. Then $\operatorname{Aff} S(G,u) = \operatorname{Aff} F \times \operatorname{Aff} F'$ (as ordered groups); let π denote the projection map to $\operatorname{Aff} F'$ (given by $\pi(a) = a|F'$). Here the kernel of π $(F)^\perp \cong \operatorname{Aff} F$ is an order ideal of $\operatorname{Aff} S(G,u)$, and admits a relative order unit (this is where the closedness of F' is used).

We claim that $\pi(J) \cap (\operatorname{Aff} F')^{++} \neq \emptyset$. As $\operatorname{Aff} F'$ is itself a dimension group, sums of order ideals are ideals, so that Lemma 1.3 applies. If the intersection is empty, then there exists a trace on $\operatorname{Aff} F'$ that kills $\pi(J)$, and this induces a trace, τ, on $\operatorname{Aff} S(G,u)$ supported on F', that kills J. But $J^\perp = Z = F$, a contradiction.

Thus there exists $a \in J$ such that $a|F' \gg 0$. For every element $j \in J$, compactness of F' implies that there exists an integer N such that $-Na|F' \leq j|F' \leq Na|F'$. On F, each of a and j restrict to zero, so that $-Na \leq j \leq Na$ (as elements of $\operatorname{Aff} S(G,u)$). Thus $j = Na - (Na - j)$ with both Na and $Na - j$ in $\operatorname{Aff} S(G,u)^+$, so that $J = P(J)$. The rest follows from [BeH, Proposition B3]. •

COROLLARY 1.4 Let (G,u) be a simple dimension group, and let U be a family of normalized traces on it. Set $Z = Z(\ker U)$. If Z is a face of $S(G,u)$ whose complementary face is closed, then $G/\ker U$ is a simple dimension group with respect to the quotient ordering.

The following is an easy corollary to a combination of earlier results and is a partial converse to [BeH, B4], in the special case that $S(G,u)$ is finite-dimensional and Z contains an interior point of $S(G,u)$. (Since Z contains an interior point, the $P(J)$ term must be zero.)

The expression *not unperforated* is cumbersome; *holey* is proposed as a synonym.

COROLLARY 1.5 Let (G,u) be an approximately divisible group such that $S(G,u)$ is finite-dimensional. Let $U \subset S(G,u)$ be a set of traces such that $Z := Z(\ker U)$ contains an interior point of $S(G,u)$. The following are equivalent.

(a) $G/\ker U$ is unperforated;

(b) the closure of the image of $\ker U$ in $\operatorname{Aff} S(G,u)$ is a vector space;

(c) for any trace τ of G, either $\tau(\ker U)$ is zero or a dense subgroup of $\mathbf{R}$;

(d) for any bounded linear functional σ on G, $\sigma(\ker U)$ is either zero or a dense subgroup of $\mathbf{R}$.

Proof. (b) implies (a) follows from Proposition 1.1. (c) implies (d). Let $z \in Z \cap \operatorname{int} S(G,u)$, and suppose that σ is a bounded linear functional such that $\sigma(\ker U) = \delta\mathbf{Z}$ (a discrete and nonzero subgroup of the reals).

Identifying $\operatorname{Aff} S(G,u)$ with $\mathbf{R}^n$, and the pure traces with the coordinate projection functions, we see that z is a strictly positive linear combination of the coordinate functions, and σ is just a linear combination of the coordinate functions. Hence there exists a positive integer N such that $\tau := \sigma + Nz$ is a positive combination of the coordinate functions, that is, an (unnormalized) trace. Obviously $\tau(\ker U) = \sigma(\ker U) = \delta\mathbf{Z}$, a contradiction.

(d) implies (b). Let J be the closure of the image of $\ker U$ in $\operatorname{Aff} S(G,u)$; as the latter is just $\mathbf{R}^n$, J decomposes as a direct sum of a (real) vector space with a discrete subgroup; the vector space is necessarily closed, so if the discrete subgroup is not zero, there exists a bounded linear functional on $\operatorname{Aff} S(G,u)$, hence on G, σ such that the range of σ on J is a discrete nonzero subgroup of the reals. It follows that $\sigma(\ker U)$ is the same, that is, a discrete nonzero subgroup, a contradiction.

(b) implies (c) is trivial. Finally, (a) implies (d) is [BeH, B4]. •

The implication (a) implies (c) is true without the finite-dimensional assumption (although its proof is rather intricate). The structure of closed subgroups of finite dimensional vector spaces is used in the proof of (d) implies (b). The proof of (c) implies (d) relies on finite-dimensionality (because even in very tractable infinite dimensional cases, z in the interior does not imply the existence of the N), but the result could conceivably be true without the finite dimensional assumption.

Suppose that $H = \ker_G Z(H)$. Let F denote the smallest face of $S(G,u)$ that contains $Z(H)$; this means that $Z(H) \subset F$ and $Z(H)$ contains a point in the relative interior of F. A minor modification of the proof of [BeH, B4] yields the following.

LEMMA 1.6 If G/H is unperforated and $H = \ker_G(Z(H))$ and F is the smallest face containing $Z(H)$, then for all traces $\tau \in F$, $\tau(H)$ is either zero or has dense image in $\mathbf{R}$.

Proof. If $Z(H) = F$, then $Z(H)$ is a face, and the conclusion trivially applies. Otherwise, $Z(H) \neq F$, and the latter is at least one-dimensional. Since F is the smallest face containing $Z(H)$, the latter contains a point z in the relative interior of F. Following the proof of [BeH, B4], there exists a line segment joining τ to z that extends outside Z, but still within F. The affine span of this segment hits the boundary of F at τ_2, yielding a decomposition $z = \lambda\tau + (1-\lambda)\tau_2$ for some some $0 < \lambda < 1$.

There exists $\delta > 0$ such that $\tau(H) = \delta\mathbf{Z}$. For any $v \in \ker_G Z(H)$, we have $z(v) = 0$, so $\tau_2(v) = -\lambda\tau(v)/(1-\lambda)$. In particular, $\tau_2(H)$ has nonzero discrete image. Now the rest of the argument of [op. cit.] applies. •

For a face F of $\operatorname{Aff} S(G,u)$, there is a natural order-preserving homomorphism $\pi_F : \operatorname{Aff} S(G,u) \to \operatorname{Aff} F$ given by $j \mapsto j|F$. A consequence of the preceding is that if G/H is unperforated, and F is the smallest face containing $Z(H)$, then $\pi_F(H)$ is dense in a (real) vector subspace of $\operatorname{Aff} F$.

LEMMA 1.7 If G/H is unperforated, $H = \ker_G(Z(H))$ and F is the smallest face containing $Z(H)$, then the closure of $\pi_F(H)$ (in $\operatorname{Aff} F$) is a vector space.

Proof. If not, from the classification of closed subgroups of Euclidean space, there exists a bounded linear functional $\sigma \in \langle F - F\rangle$ such that $\sigma(\pi_F(H)) = \delta\mathbf{Z}$ for some nonzero δ. Via π_F, we can regard σ as a bounded linear functional that can be written in the form $\sigma = \sum_{\tau_i \in F} a_i\tau_i$ with a_i real. Since F is the smallest face containing $Z(H)$, the latter contains a point z in the relative interior, hence we can write $z = \sum_{\tau_i \in F} b_i\tau_i$ where all $b_i > 0$. Hence there exists N such that $Nz + \sigma$ is a trace, and its normalized form will thus be an element of F whose image is properly cyclic. This contradicts the preceding. •

In Example 1, if $\{w_i\}$ is linearly independent over the reals, then it is immediate that $Z(H) = (\tau_1+\tau_2)/2$, and then F is just the edge joining τ_1 to τ_2.

If $A = \mathrm{I}$ and $\{w_i\}$ is not linearly independent and all the $\alpha_i > 0$, then it is routine to see that $\rho := \tau_2 + \sum \alpha_i\tau_{i+2}$ kills H, and in fact $Z(H)$ consists of the edge joining $(\tau_1+\tau_2)/2$ to the normalized form of ρ, $\rho/(1+\sum\alpha_i)$. The resulting $Z(H)$ thus contains an interior point of the standard simplex.

In all cases, we can wrap a simple dimension group G around H (so $H \subset G \subset \mathbf{R}^n$, G is dense in $\mathbf{R}^n$), and do so so that $H = \ker_G H$ (Lemma 1.12).

A necessary condition for G/H to be unperforated in these examples when $Z(H)$ is a singleton) is merely that all traces in the face generated by τ_1 and τ_2 have either zero or dense image; since the average kills H, the condition is satisfied because of our hypotheses, namely that $\sum\alpha_i\mathbf{Z}$ has rank at least two.

On the other hand, if $\{w_i\}$ is not linearly independent, then $Z(H)$ contains an interior point, and thus $F = S(G,u)$. Then the necessary condition that G/H be unperforated is more drastic: that the closure of H within $\mathbf{R}^n$ be a vector space. This forces $\{1, \eta_1, \dots, \eta_{n-2}\}$ to be rationally linearly independent. We will deal with Example 1 less casually soon.

The condition given in Lemmas 1.6 or 1.7 is relatively easy to verify; the natural question is whether the converse holds. We will prove that this is the case.

From now until further notice, we assume G is a simple dimension group with finitely many pure traces.

LEMMA 1.8 Suppose H is a subgroup of the dense subgroup G of $\mathbf{R}^n$ such that $H \cap G^{++} = \emptyset$ and G/H is torsion-free. Sufficient for G/H to be unperforated is that the closure of $\pi_F(H)$ is a vector space, and that for all pairs (ϵ, N), there exists $j \in H$ such that $\widehat{j} \geq -\epsilon$ and $j|F' > N$.

Proof. Pick $h \in H$ and $h_1 \in H$ such that $\left\| \widehat{2h - h_1|F} \right\| < \epsilon/2$. Select $j \in H$ such that $\widehat{j} \geq -\epsilon/4$ and $\widehat{j}|F' > |\min\left(\widehat{h} - \widehat{h_1}/2\right)|F'|$. Then $h - 2h_1 + 2j \geq -\epsilon$ (since this occurs on both F and F'), and thus $h - 2(h_1 - j)$ has the requisite property. •

LEMMA 1.9 Suppose $H = \ker{}_G Z(H)$, where G is a dense subgroup of $\mathbf{R}^n$, equipped with the strict ordering, and F is the smallest face containing $Z(H)$. Then $F^\perp \cap (H \cdot \mathbf{R})$ contains an element j such that $\pi_{F'}(j)$ is strictly positive.

Remark. The subscript on the last line of the statement of the lemma reads F', not F.

Proof. We verify (*) of Proposition 1.1. If $F = S(G,u)$, F' is empty, and there is nothing to do. Otherwise, it is convenient to relabel the vertices of F as $\{\tau_1, \dots, \tau_k\}$ and those of F' as $\{\tau_{k+1}, \dots, \tau_n\}$. We want to show there exists $j \in H \cdot \mathbf{R}$ of the form $(0, 0, \dots, 0; *, *, \dots, *)$ where the asterisks represent strictly positive real numbers, and the initial k entries are zero.

Let $I = F^\perp \cap (H \cdot \mathbf{R})$ (consisting of all the elements of $H \cdot \mathbf{R}$ whose initial k terms are zero). Suppose there exists a trace $\rho \in F'$ such that $\rho(I) = \{0\}$. (Any $\sigma \in F$ will kill automatically kill I.) Then ρ induces a linear functional $\phi : \pi_F(H \cdot \mathbf{R}) \to \mathbf{R}$, via $\pi_F(H \cdot \mathbf{R}) \cong (H \cdot \mathbf{R})/I$. We may extend ϕ to a linear functional on $\pi_F(\mathbf{R}^n)$, and then lift it to a linear functional on $\mathbf{R}^n$ which factors through π_F; call the resulting linear functional ϕ as well.

Then $\phi = \sum_{i=1}^k r_i \tau_i$ is supported on the vertices of F (since ϕ factors through π_F) and $\phi|H = \rho|H$. By hypothesis, there exists $z \in Z(H)$ and in the relative interior of F—hence there exists $N > 0$ such that $\psi := \phi - Nz$ is a negative linear functional (supported on F with all negative coefficients) and $\psi|H = \rho|H$. Then $\rho - \psi$ is a positive linear functional that kills H. Since $Z(H) \subset F$ and the normalized form of $-\psi$ is in F, this forces $\rho \in F$, a contradiction.

Thus for all $\rho \in F'$, $\rho(I) \neq \{0\}$. This means that the image of I, this time under $\pi_{F'}$, call it $J \subset \mathbf{R}^{\{k+1,\dots,n\}}$ is not killed by any trace of the latter. By Lemma 1.3, J contains a strictly positive element, and this implies I has an element whose coordinates beyond the kth are strictly positive. •

The l^2-norm is used in the following well-known result.

LEMMA 1.10 Let H be a full lattice in Euclidean space V. Given $v \in V$, for all $\epsilon > 0$, for all real $R > 0$, there exists an integer $r > R$ and $h \in H$ such that $\|h - rv\| < \epsilon$.

Proof. Fix $\epsilon > 0$ and $R > 0$. The set

$$\mathcal{C} := \{x \in V \mid \|x - \langle x, v\rangle v\| < \epsilon\}$$

is convex and 0-symmetric (meaning that the set of rational points therein is stable with respect to multiplication by -1), with infinite volume. Thus, by a theorem of van der Corput ([vdC] or [GL, 7.2]), it contains infinitely many points of H. In particular, it contains $\pm h$ with $h \in H$ having norm $\|h\| > \epsilon + R\|v\|$. Choosing h so that $r := \langle h, v\rangle$ is positive, we conclude that the pair (r, h) has the required property. •

THEOREM 1.11 Let G be a simple noncyclic dimension group with finitely many pure traces, and let H be a subgroup of G of the form $\ker{}_G Z(H)$. Let F be the smallest face containing $Z(H)$. Then G/H is unperforated iff for all $f \in F$, $f(H)$ is either zero or dense in $\mathbf{R}$.

Proof. One direction (which does not require the finite-dimensional hypothesis) is Lemma 1.6. For the other, assume that $f(H)$ is zero or dense for each $f \in F$. Then $\pi_F(H)$ is dense in $\mathrm{Aff}\, F$. Form $W = H \cdot \mathbf{R}$, the vector space generated by H, and define $W^+ = W \cap (\mathbf{R}^n)^+$ (identifying $\mathrm{Aff}\, S(G,u)$ with $\mathbf{R}^n$).

Then we have $Z(H) = Z(W)$. Index the pure traces $\{\tau_i\}$ so that $\tau_i \in F$ iff $i \leq k$. If $k = n$, that is, $F = S(G,u)$, then $Z(H)$ contains an interior point, and the result follows from Corollary 1.5. Otherwise, $k < n$, and $F^\perp$ consists precisely (after our relabelling of the indices) of all elements of $\mathbf{R}^n$ of the form $(0^k, w)$ where w is an arbitrary $n-k$-tuple. In particular, W^+ contains all the elements of this form where $w \gg 1$ entrywise; let $v = (0^k, w)$. By Lemma 1.9, such an element exists in $H \cdot \mathbf{R}$.

Inside H, we may find an $\mathbf{R}$-basis for W, and let H_0 be the lattice (discrete subgroup) the basis generates. Select $\epsilon, N > 0$. By Lemma 1.10, there exists a positive integer c such that $c > N$ and $h \in H_0 \subset H$such that $\|h - cv\| < \epsilon$. Since $v|F$ is zero, we have that $\|h|F\| < \epsilon$, while $h|F' > c - \epsilon > N$. This verifies the criterion in Lemma 1.8. •

Theorem 1.11 is a result about subgroups of dimension groups. For the purposes of Diophantine inequalities, we can remove the existence of G from the statement and rephrase it entirely in terms of H, provided we view the latter as a subgroup of $\mathbf{R}^n$. We of course require H to contain no strictly positive elements, so that by Lemma 1.3, $Z(H)$ (computed with respect to this embedding of H in $\mathbf{R}^n$) is not empty. This permits us to define F, the smallest face of the standard simplex that contains $Z(H)$, without reference to G.

LEMMA 1.12 Let H be a subgroup of $\mathbf{R}^n$ such that $H \cap (\mathbf{R}^n)^{++} = \emptyset$. Then there exists a subgroup G of $\mathbf{R}^n$ with the following properties.

(i) $H \subset G$ and $u = (1, 1, \dots, 1) \in G$;
(ii) G is dense in $\mathbf{R}^n$;
(iii) $\ker_G Z(H) = H$
(iv) G/H is free and $\mathrm{rank}\, G/H = n - \dim H \cdot \mathbf{R} + 1$.

Proof. First, we show that $u \notin H \cdot \mathbf{R}$. Otherwise, we could find a finite subset $\{h_i\}$ of H, together with a finite set of real numbers r_i such that $\sum h_i r_i = u$. Approximate each r_i by rational numbers, $q_{j,i}$ (so $\{q_{j,i}\} \to r_i$ for each i), and set $s_j = \sum h_i q_{j,i}$. Then $s_j \to u$; since u is strictly positive, there exists k such that s_k is strictly positive. As the coefficients are rational, there exists an integer N such that $N q_{k,i}$ are integers for all i. But then $N s_k$ is a strictly positive element of H, a contradiction.

Fix $\tau \in Z(H)$ ($Z(H)$ is nonempty by Lemma 1.3).

Now choose a basis for $H \cdot \mathbf{R}$ consisting of elements of H, say $B := \left\{h^1, \dots, h^m\right\}$. By the preceding, $B' = B \cup \{u\}$ is linearly independent over $\mathbf{R}$, and thus we may complete B' to a real basis for $\mathbf{R}^n$, $B'' = B' \cup \{g_{m+2}, \dots, g_n\}$ with the additional property that $\tau(g_j) \neq 0$ (this can always be arranged). Pick a set of real numbers θ_i such that the set $\{1, \theta_1, \dots, \theta_n\}$ is rationally linearly independent. Form the field K generated over the rationals by $\{\theta_{m+1}, \dots, \theta_n\}$ and let $\{e_{m+2}, \dots, e_n\}$ be a subset of the reals such that $\{1, e_{m+2}, \dots, e_n\}$ is linearly independent over K.

Set $b_{m+1} = u$ and $b_{m+i} = g_{m+i} \cdot e_{m+i}/\tau(g_{m+i})$ if $i > 1$. Then $B''' := B \cup \{b_{m+i}\}_{i \geq 1}$ is obviously a basis. Define $b_{n+1} = \sum h_i \theta_i + \sum_{j=m+1}^n b_j \theta_j$. Finally, we set $G = H + \sum_{j=m+1}^{n+1} b_j \mathbf{Z}$. Obviously (i) holds. Moreover, G contains the set $\left\{h^1, h^2, \dots, h^m; b_{m+1}, \dots, b_n\right\}$, a basis for $\mathbf{R}^n$, together with the element b_{n+1}, and of course they generate a dense (free of rank $n+1$) subgroup of $\mathbf{R}^n$. So (ii) holds.

To verify (iii), suppose that $g = h + \sum_{j=m+1}^n c(j) b_j + c b_{n+1}$ is in $\ker_G Z(H) = Z(H)^{\vdash} \cap G$, where $c(j), c$ are all integers and $h \in H$. Applying $\tau \in Z(H)$, we have (since $\tau(H \cdot \mathbf{R}) = 0$)

$$
\begin{aligned}
0 &= \sum_{j=m+1}^{n} c(j)\tau(b_j) + c\tau(b_{n+1}) \\
&= \sum_{j=m+1}^{n} \tau(b_j)(c(j) + c\theta_j) \\
&= 1 \cdot (c(m+1) + c\theta_{m+1}) + \sum_{j=m+2}^{n} e_j \cdot (c(j) + c\theta_j).
\end{aligned}
$$

Since all the coefficients belong to K, linear independence over K of $\{1, e_{m+2}, \ldots, e_n\}$ forces all the coefficients to be zero, that is, $c(j) + c\theta_j = 0$ for all $m + 1 \leq j \leq n$. Linear independence over the rationals of the $\{1, \theta_1, \ldots, \theta_n\}$ forces $c(j) = c = 0$ for all j, and thus $g = h$.

Property (iv) is obvious. •

This permits us to restate the criterion without reference to dimension groups. The presence of $u = (1, 1, \ldots, 1)$ means that when we take (G, u) as obtained in Lemma 1.12, the affine representation restricts to the original embedding of $H \subset \mathbf{R}^n$.

COROLLARY 1.13 Let H be a subgroup of $\mathbf{R}^n$ containing no strictly positive elements. Then H satisfies (*) (the systems of linear inequalities can always be solved) iff for all $f \in F$, either $f(H) = 0$ or $f(H)$ is dense in $\mathbf{R}$, where F is the smallest face in the standard simplex, S, containing $Z(H) = \{\tau \in S \mid \tau(H) = 0\}$.

It need not be the case that $H \cdot \mathbf{R} = Z(H)^{\perp}$, as in the following example. First, we name the condition appearing in Lemma 1.8.

$$\text{For all } \epsilon, N, \text{ there exists an element of } H \cdot \mathbf{R},\ f, \text{ such that } f|F' > N \text{ and } \|f|F\| < \epsilon. \qquad (**)$$

If F' is empty, we disregard the relation involving N.

The following family of examples (indexed by the real parameter r) illustrates a number of points.

EXAMPLE 2 Let $\{\alpha, r, r_2, r_3, \gamma_1, \ldots, \gamma_4\}$ be an algebraically independent (over the rationals) set of positive real numbers. Define the five elements of $\mathbf{R}^{1\times 4}$ (rows of size four),

$$\begin{array}{rcl} v_1 & = & (1 \quad -1 \quad 1 \quad 0) \\ v_2 & = & (\alpha \quad -\alpha \quad r \quad 1) \\ v_3 & = & (r_2 \quad 0 \quad 0 \quad 0) \\ v_4 & = & (0 \quad r_3 \quad 0 \quad 0) \\ v_5 & = & (\gamma_1 \quad \gamma_2 \quad \gamma_3 \quad \gamma_4). \end{array}$$

Let G be the subgroup of $\mathbf{R}^{1\times 4}$ generated by $\{v_1, \ldots, v_5\}$. The determinant of the matrix consisting of the top four rows is $r_2 r_3 \neq 0$, so $\{v_1, v_2, v_3, v_4\}$ is linearly independent over the reals. This means that we can write $v_5 = \sum_{i=1}^{4} t_i v_i$ for some $t_i \in \mathbf{R}$, and the algebraic independence assumption more than yields that $\{1, t_1, t_2, t_3, t_4\}$ is rationally linearly independent. Hence G is a dense rank five subgroup of $\mathbf{R}^4$.

Let τ_i, $i = 1, 2, 3, 4$ denote the four coordinate projections, and set $\tau = \tau_1 + \tau_2$. Then it again follows from the algebraic independence that $\ker_G \tau := \tau^{\perp} \cap G = \langle v_1, v_2 \rangle$. In particular, we see that $\tau^{\perp}$ does not contain a basis for $\ker \tau$.

Define $H = \langle v_1, v_2 \rangle$; then $Z(H) = \{\tau/2\}$ when $\alpha < r$, and

$$Z(H) = \operatorname{cvx}\{\tau/2, (\tau_2 + \tau_3 - (r - \alpha)\tau_4)/(2 - r + \alpha)\}$$

(a line segment) when $\alpha \geq r$. When $\alpha > r$, H is also killed by the non-positive linear functional, $\phi := \tau_1 - \tau_3 + (r - \alpha)\tau_4$, and in fact, the annihilator of H (in the dual space spanned by the τ_i) is spanned by $\{\phi, \tau\}$. In particular, $H = \ker_G Z(H)$ in this case, and it is elementary that $H = \ker_G Z(H)$ regardless of the value of r.

Fix r. Assume (**) holds. There exist integers A, B such that $Av_1 + Bv_2 \gg -(\epsilon, -\epsilon, N, N)$. This yields $|A + B\alpha| < \epsilon$ (from the first two entries), and $A + Br > N$, and $B > N$. In particular, B has to be large and positive. If $r < \alpha$, then

$$N < A + Br = A + B\alpha + B(r - \alpha) < \epsilon - B|r - \alpha| < \epsilon,$$

a contradiction. Hence (**) fails if $r < \alpha$.

If $r > \alpha$, then $A + Br > -\epsilon + B(\alpha - r)$, and as $\epsilon \to 0$, $|B| \to \infty$, so we can always replace A, B by their negatives if necessary, and thus (**) succeeds.

When $r > \alpha$, (**) succeeds, so (*) occurs. On the other hand, we can show by a similar easy computation that (*) fails when $r < \alpha$. We show that $2h - v_1 \geq -2\epsilon(1, 1, 1, 1)$ cannot be solved for $h \in H$, provided ϵ is sufficiently small. Suppose to the contrary that $h = Av_1 + B_v$ exists; then we deduce the equations

$|2A+\alpha(2B-1)|<2\epsilon$, $2A-r(2B-1)\geq -2\epsilon$, and $r(2B-1)\geq -2\epsilon$. Since $r>0$, the last entails $B\geq 1$ (provided $\epsilon<1/2, r/2$). Then, almost identical to the previous computation,

$$2A-r(2B-1)=2A-\alpha(2B-1)-(\alpha-r)(2B-1)<2\epsilon-(2B-1)|\alpha-r|;$$

as $\epsilon\to 0$, $|B|\to\infty$, so $2B-1\to\infty$ (since B is already known to be positive), so the right side goes to $-\infty$, a contradiction. [Essentially the same result occurs if we try to halve v_2 rather than v_1.]

When $r>\alpha$, $\ker Z(H)$ is three-dimensional, hence strictly contains $H\cdot\mathbf{R}$. It is easy to see that $\ker Z(H)$ contains lots of elements which are strictly positive on F', but we have to find one in $H\cdot\mathbf{R}$ to prove (*) holds. •

Corollary 1.13 allows us to give quick answers as to which cases of Example 1 (in the Introduction) satisfy (*). For a real matrix C, let $\mathcal{R}(C)$ denote its row space.

EXAMPLE 1 (REVISITED) Consider the real matrix,

$$B=\begin{pmatrix} \alpha_1 & -\alpha_1 & & & & & & \\ \alpha_2 & -\alpha_2 & & & & & & \\ \vdots & \vdots & & & A & & & \\ \alpha_{n-2} & -\alpha_{n-2} & & & & & & \\ \alpha_{n-1} & -\alpha_{n-1} & \eta_1 & \eta_2 & \eta_3 & \dots & \eta_{n-3} & \eta_{n-2} \end{pmatrix}.$$

Assume that at least one ratio α_i/α_j is defined and irrational. Let H be the abelian group generated by the rows of B. Let B_+ and B_- denote the matrices obtained by deleting the second, respectively first column of B. Let $\tau=(\tau_1+\tau_2)/2\in Z(H)$.

(0) If $Z(H)=\{\tau\}$, then H satisfies (*).

(i) $Z(H)=\{\tau\}$ iff both $\mathcal{R}(B_+)\cap(\mathbf{R}^{n-1})^{++}\neq\emptyset$ and $\mathcal{R}(B_-)\cap(\mathbf{R}^{n-1})^{++}\neq\emptyset$.

(ii) If the rows of B form a real linearly independent set, that is, $\operatorname{rank}B=n-1$, then $Z(H)=\{\tau\}$ and H satisfies (*).

(iii) Suppose that $A=\mathrm{I}$, $\operatorname{rank}B=n-2$, and $\alpha_2\neq 0$.
  (a) If not all nonzero α_i $(i\leq n-2)$ have the same sign, then $Z(H)=\{\tau\}$ and H satisfies (*);
  (b) If all the nonzero α_i have the same sign, then H satisfies (*) iff $\{1\}\cup\{\eta_j\}_{j\in S}$ is rationally linearly independent, where $S=\{j\leq n-2\mid \alpha_j\neq 0\}$; in these cases, $Z(H)$ is one-dimensional (a line segment).

Proof. Since $(\tau_1+\tau_2)/2\in Z(H)$, the smallest face containing $Z(H)$ contains the convex hull of $\{\tau_1,\tau_2\}$, with equality if $Z(H)=\{\tau\}$.

(0) An immediate consequence of Theorem 1.11; any trace in $F=\operatorname{cvx}\{\tau_1,\tau_2\}$ is congruent to a multiple of τ_1 when restricted to H, so its range is dense.

(i) Let $\sigma=\sum a_i\tau_i$ be a positive linear functional that kills H, and at least one of $a_i>0$ for $i\geq 3$. Since $\tau\in Z(H)$, we may assume that at least one of $a_1=0$ or $a_2=0$ holds. Define the columns $a_+=(\alpha_1,\alpha_3,\dots\alpha_n)^T$ and $a_-=(\alpha_2,\alpha_3,\dots\alpha_n)^T$, so that $\sigma(H)=0$ is equivalent to either $B_+a_+=\mathbf{0}$ or $B_-a_-=\mathbf{0}$. Since $a_\pm$ are both nonzero (as $a_i\neq 0$ for at least one $i\geq 3$) and nonnegative, Gordan's Theorem applies, and we see that at least one of $B_\pm\boldsymbol{x}=\mathbf{0}$ admits a nonnegative nonzero solution iff one of $\mathcal{R}(B_\pm)\cap(\mathbf{R}^{n-1})^{++}=\emptyset$.

(ii) Since the rows are linearly independent, $\dim H\cdot R=n-1$, so the annihilator of H must be one dimensional, hence its intersection with $S(G,u)$ can only be a point or empty. But we already have that $\tau\in Z(H)$, whence $Z(H)=\{\tau\}$.

(iii) Since $\operatorname{rank}B=n-2$, $\dim H\cdot\mathbf{R}=n-2$, so its annihilator is two-dimensional. We see that $\phi=\tau_2+\sum_{i\geq 3}\alpha_{i-2}\tau_i$ kills H; hence every linear functional that kills H is of the form $\psi=r\tau+s\phi$.

If at least two of the α_i have different signs, if such a ψ were a positive linear functional, then $s=0$; hence when the signs are different, $Z(H)=\{\tau\}$, which together with (0) yields (a).

If all the signs of the α_i are equal, then it is immediate that there is a positive linear functional of the form $\rho=r\tau+s\phi$. We have that either $\phi=\tau_2+\sum_{j\in S}\alpha_j\tau_{j+2}$ or its negative is a trace killing H, so the smallest face, F, containing $Z(H)$ is $\operatorname{cvx}\{\tau_1,\tau_2\}\cup\{\tau_{j+2}\}_{j\in S}$. By Lemma 1.7, Theorem 1.11, and Corollary 1.13, H satisfies (*) iff the closure of $\pi_F(H)$ in $\operatorname{Aff}F$ is a vector space. Since w_{n-1} is a real linear combination of $\{w_i\}_{i \leq n-2}$, it must be that $w_{n-1} = \sum \eta_i w_i$. As $\{\pi_F(w_j)\}_{j \in S}$ is linearly independent, the closure of $\pi_F(H)$ is a real vector space iff iff $\{1\} \cup \{\eta_j\}_{j \in S}$ is rationally linearly independent. •

Refinability

A good reason to consider unperforation of G/H comes from the study of refinable traces (originally refinable measures on Cantor dynamical systems, [Ak]). A trace τ on a partially ordered abelian group (G, u) is *refinable* if for $g \in G^+ \setminus \{0\}$ and for finite collections $\{a_i\} \subset G^+$ such that $\sum \tau(a_i) = \tau(g)$, there exist $a_i' \in G^+$ such that $\tau(a_i) = \tau(a_i')$ and $g = \sum a_i$. The trace τ is *good* if for $a \in G$, $g \in G^+ \setminus \{0\}$ with $\tau(a) < \tau(g)$, there exists $a' \in G^+$ such that $\tau(a) = \tau(a')$ and $a \leq g$.

There are generalizations, variations, and implications discussed in [BeH]. Good traces are refinable, but not conversely. When G is a simple dimension group, a trace τ is good iff the closure of the image of $\ker \tau$ in $\operatorname{Aff} S(G, u)$ is $\tau^{\vdash} = \{h \in \operatorname{Aff} S(G, u) \mid h(\tau) = 0\}$ [BeH, 1.7], and this characterization is very useful. Left open in op.cit., was a corresponding characterization of refinability for traces, because it was not clear that $G/\ker \tau$ was unperforated. This can now be addressed.

To complete the characterization, we need yet another definition from [BeH]. Let $U \subseteq S(G, u)$; we say that U is *good with respect to* $S(G, u)$ if whenever $a, b \in \operatorname{Aff} S(G, u)^+$ are such that $a|U \ll b|U$, then there exists $a' \in \operatorname{Aff} S(G, u)^+$ such that $a'|U = a|U$ and $a' \ll b$. When U is an intersection of a closed flat with $S(G, u)$ (for example, it could be of the form $Z(H)$ for H a subgroup of $\operatorname{Aff} S(G, u)$), exactly which such sets are good is completely determined—in the finite-dimensional case [H2, B.3]; they are given as coproducts (of a special type) in the family of simplices with subsimplices.

COROLLARY 2.1 Suppose G is a simple dimension group and $S(G, u)$ is finite-dimensional. Then a trace τ is refinable iff

(a) $G/\ker \tau$ is unperforated

(b) $Z(\ker \tau)$ is good with respect to $S(G, u)$

(c) the closure of $\ker \tau$ in the affine representation is $Z(H)^{\vdash}$.

Proof. We have $\ker \tau = \ker (Z(\ker \tau))$. If τ is refinable, then for every trace σ, $\sigma(\ker \tau)$ is not properly cyclic [BeH, B5], and by Theorem 1.11, it follows that $G/\ker \tau$ is unperforated. The rest of the both sets of implications follow from [BeH, 7.6]. •

COROLLARY 2.2 Hypotheses as in the previous. The following are equivalent.

(i) τ is good

(ii) τ is refinable and $Z(\ker \tau) = \{\tau\}$.

Recall from [H], that a *critical dimension group of rank* $n + 1$ is a dense subgroup of $\mathbf{R}^n$, free of rank $n + 1$, equipped with the strict ordering inherited from $\mathbf{R}^n$ (that it is a dimension group is a consequence of [EHS]). If H is a subgroup of lesser rank of a critical dimension group, then H is discrete.

As in [BeH], sufficient for a trace τ to be refinable is that $\ker \tau = \operatorname{Inf} G$; the set of such traces is a dense G_δ of the normalized trace space [GHH], and in general the set of refinable traces contains a dense G_δ.

When all subgroups of rank less than full are discrete, the only refinable traces are those with zero kernel, so at least in that case, they form a dense G_δ.

COROLLARY 2.3 Let G be a critical dimension group of rank $n + 1$, with $n \geq 2$. Then a trace τ is refinable iff $\ker \tau = \{0\}$.

Proof. Let $H = \ker \tau$. By [BeH, B3], for every trace σ, $\sigma(H)$ is not properly discrete, so 1.11 applies, so that G/H is unperforated. Then [BeH, 7.6e] implies that the closure of H in $\mathbf{R}^n$ is a real vector space. However, H is discrete, being of rank at most $n - 1$. •

Appendix. Gordan's theorem and Farkas' lemma

Gordan's theorem (not to be confused with Gordan's lemma, a result concerning toric varieties) asserts the following.

THEOREM A1 (Gordan) Let A be an $m \times n$ real matrix. Exactly one of the following is true.

(1) There exists $Y \in (\mathbf{R}^{n \times 1})^+ \setminus \{\mathbf{0}\}$ such that $AY = \mathbf{0}$.

(2) There exists $X \in \mathbf{R}^{1 \times m}$ such that XA is strictly positive.

If (2) holds, then we may find X to have only integer entries, by the obvious density argument. Then Lemma 1.3 can be seen as a generalization of Gordan's Theorem. Set $G = \mathbf{R}^{n\times 1}$ with the usual coordinatewise ordering, let L be the subgroup of G generated by the rows of A. The traces of G are given by right multiplication by columns of nonnegative vectors, so alternative (1) says there exists a trace τ such that $\tau(H)$ is zero. Alternative (2) with $X \in \mathbf{Z}^{1\times m}$ says that $\sum x_i w_i$ (where w_i are the rows of A) is strictly positive, hence an order unit of H.

Farkas' lemma does not generalize terribly well, partly because of the possible presence of infinitesimals, but for other reasons as well. There are numerous equivalent forms; that presented here is the transpose of one given in [Wi].

THEOREM A2 (Farkas) Let A be a real $m \times n$ matrix, and let $b \in \mathbf{R}^{1\times n}$. Exactly one of the following is true.

(1) There exists $X \in \mathbf{R}^{1\times m}$ such that $XA \leq b$.

(2) There exists nonzero $Y \in (\mathbf{R}^{n\times 1})^+$ such that $AY = \mathbf{0}$ and $bY < 0$.

To attempt to generalize this, let (G, u) be a partially ordered abelian group, let H be a subgroup, and let $g \in G$. Then we *hope* that either (1) there exists $h \in H$ such that $h \leq g$ or (2) there exists a trace τ of G such that $\tau(H) = 0$ and $\tau(g) < 0$.

If we take $G = \mathbf{R}^{1\times n}$ (usual ordering), $g = b$, and H the group generated by the rows of A, then right multiplication by the nonnegative nonzero Y amounts to taking a trace. There is a slight problem, in that in the matrix formulation X is allowed to vary over the real rows, while in the ordered group, they are restricted to the integer entries; this didn't present a problem in the case of Gordan's theorem, but it explains why we are going to need at least unperforated in the hypotheses to generalize Farkas' lemma.

So suppose we try to prove this ordered group version of Farkas' lemma. If (1) fails, the condition becomes $(g + H) \cap G^+ = \emptyset$—in other words, $g + H$ is not in $(G/H)^+$ (assuming that the pre-ordering on G/H is a partial ordering).

We have an easy lemma.

LEMMA A3 Let (G, u) be a partially ordered abelian group, and let H be a subgroup of G such that G/H is partially ordered (with respect to the quotient ordering). Suppose $g \in G$ such that $(g + H) \cap G^+ = \emptyset$. If G/H is unperforated, there exists $\tau \in S(G, u)$ such that $\tau(g) \leq 0$ and $\tau|H \equiv 0$.

Remark. Aside from unperforation and partial ordering hypotheses on G/H (the latter is convexity of H, that is, if $h_1 \leq j \leq h_2$ with $h_i \in H$ and $j \in G$, then $j \in H$), this result is weak, as it only gives $\tau(g) \leq 0$, not the desired $\tau(g) < 0$.

Proof. We have $g + H \notin (G/H)^+$. (i) As G/H is unperforated, there exists a trace σ on $(G/H, u + H)$ such that $\sigma(g + H) \leq 0$. Then σ composed with the quotient map is the desired τ. •

Of course, if we knew that G/H was archimedean (in the usual sense), then the same argument would yield a trace with $\tau(g) < 0$ and $\tau|H \equiv 0$.

In the real case, that is, where we take the row space for H rather than the abelian group generated by the set of rows, both G and H will be divisible, hence G/H is automatically unperforated.

References

[Ak] E Akin, *Measures on Cantor space,* Topology Proc, 24 (1999) 1–34.

[BeH] S Bezuglyi & D Handelman, *Measures on Cantor sets: the good, the ugly, the bad,* Trans Amer Math Soc 366 (2014) 6247–6311.

[vdC] JG van der Corput, *Verallgemeinerung einer Mordellschen Beweismethode in der Geometrie der Zahlen,* Acta Arith 1 (1935), 62–66.

[EHS] EG Effros, David Handelman, & Chao-Liang Shen, *Dimension groups and their affine representations,* Amer J Math 102 (1980) 385–407.

[GHH] T Giordano, D Handelman, M Hosseini, *Orbit equivalence of Cantor minimal systems and their continuous spectra,* in preparation.

[G] KR Goodearl, *Partially ordered abelian groups with interpolation,* Mathematical Surveys and Monographs, 20, American Mathematical Society, Providence RI, 1986.

[GH1] KR Goodearl & D Handelman, *Rank functions and K_0 of regular rings,* J Pure and Applied Alg 7 (1976) 195–216.

[GH2] KR Goodearl & David Handelman, *Metric completions of partially ordered abelian groups,* Indiana Univ J Math 29 (1980) 861–895.

[GL] PM Gruber & CG Lekkerkerker, *Geometry of numbers,* North Holland, New York 1987.

[H] D Handelman, *Free rank $n+1$ dense subgroups of $\mathbf{R}^n$ and their endomorphisms,* J Funct Analysis 46 (1982), no. 1, 1–27.

[H2] D Handelman, *Good measures for non-simple dimension groups, Arχiv,*

[Wi] Wikipedia, *Wikipedia article on Farkas' lemma,* http://en.wikipedia.org/wiki/Farkas'_lemma

Mathematics Department, University of Ottawa, Ottawa ON K1N 6N5 Canada;
dehsg@uottawa.ca (dh) & droy@uottawa.ca (dr).